

Dimetapp Multi-Symptom Cold & Flu

Drug Facts

Active ingredients (in each 10 mL)

Acetaminophen, 320 mg	Pain reliever/fever reducer
Diphenhydramine HCl, 12.5 mg	Antihistamine/cough suppressant
Phenylephrine HCl, 5 mg	Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - nasal congestion
 - cough
 - itching of the nose or throat
 - temporarily reduces fever
 - temporarily restores freer breathing through the nose
- headache
- sore throat
- runny nose
- sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 6 doses in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to sedate a child or to make a child sleep
- in a child under 6 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin

Drug Facts (continued)

Ask a doctor before use if user has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if user is

- taking the blood thinning drug warfarin
- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occurs
- pain, cough, or nasal congestion gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions

- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

Drug Facts (continued)

age	dose
adults and children 12 years and over	20 mL every 4 hours do not take more than 6 doses in any 24-hour period
children 6 to 12 years	10 mL every 4 hours do not give more than 5 doses in any 24-hour period
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 7 mg and potassium 6 mg
- store at 20-25°C (68-77°F)
- read and keep carton

Inactive ingredients anhydrous citric acid, artificial flavor, edetate disodium, FD&C red no. 40, glycerin, menthol, polyethylene glycol, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? Call 1-888-594-0828 weekdays 9 AM to 5 PM EST