

# Dimetapp Nighttime Cold & Congestion

## Drug Facts

### Active ingredients (in each 10 mL)

Diphenhydramine HCl, 12.5 mg .....	Antihistamine/ cough suppressant
Phenylephrine HCl, 5 mg .....	Nasal decongestant

### Purposes

### Uses

- temporarily relieves:
  - nasal congestion
  - runny nose
  - cough
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily restores freer breathing through the nose

### Warnings

#### Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other product containing diphenhydramine, even one used on skin

#### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

#### Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

## Drug Facts (continued)

### When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

### Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occurs
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

### Other information

- each 10 mL contains: sodium 8 mg
- store at 20-25°C (68-77°F) ■ do not refrigerate

**Inactive ingredients** anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

**Questions or comments?** Call 1-888-594-0828 weekdays 9 AM to 5 PM EST