Dimetapp Cold & Allergy

Drug Facts Active ingredients

(in each 10 mL) Purposes Brompheniramine maleate, 2 mg...... Antihistamine Phenylephrine HCl, 5 mg...... Nasal decongestant Uses temporarily relieves: nasal congestion
 runny nose sneezing itching of the nose or throat itchy, watery eyes due to hay fever temporarily restores freer breathing through the nose Warnings Do not use to sedate a child or to make a child sleepy
 if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions,

- or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate
- gland gland a breathing problem such as emphysema, asthma, or chronic bronchitis
- Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or
- stimulant ٠
- taking sedatives or tranquilizers

Drug Facts (continued) When using this product

- do not use more than directed
 drowsiness may occur
- avoid alcoholic beverages
 alcohol, sedatives, and tranquilizers may
- increase drowsiness be careful when driving a motor vehicle or
- operating machinery excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occurs
 symptoms do not get better within 7 days or are
- accompanied by fever

If pregnant or breast-feeding, ask a health

professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

do not take more than 6 doses in any 24-hour period

measure only with dosage cup provided
 keep dosage cup with product
 measure only with product

age	dose	
adults and children 12 years and over	20 mL every 4 hours	
children 6 to under 12 years	10 mL every 4 hours	
children under 6 years	do not use	

Other information

 each 10 mL contains: sodium 6 mg
 store at 20-25°C (68-77°F) Inactive ingredients anhydrous citric acid. artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? Call 1-888-594-0828 weekdays 9 AM to 5 PM EST