

Dimetapp Cold & Cough

<h2>Drug Facts</h2>	
<p>Active ingredients (in each 10 mL)</p> <p>Brompheniramine maleate, 2 mg.....Antihistamine Dextromethorphan HBr, 10 mgCough suppressant</p>	<p>Purposes</p>
<p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves: <ul style="list-style-type: none"> ■ cough ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes due to hay fever (or other upper respiratory allergies) ■ temporarily alleviates the intensity of coughing 	
<p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ to sedate a child or to make a child sleepy ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ trouble urinating due to an enlarged prostate gland ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema <p>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ do not use more than directed ■ may cause marked drowsiness ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children 	

<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>								
<p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than 6 doses in any 24-hour period ■ measure only with dosage cup provided ■ keep dosage cup with product ■ mL = milliliter <table border="1"> <thead> <tr> <th>age</th> <th>dose</th> </tr> </thead> <tbody> <tr> <td>adults and children 12 years and over</td> <td>20 mL every 4 hours</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>10 mL every 4 hours</td> </tr> <tr> <td>children under 6 years</td> <td>do not use</td> </tr> </tbody> </table>	age	dose	adults and children 12 years and over	20 mL every 4 hours	children 6 to under 12 years	10 mL every 4 hours	children under 6 years	do not use
age	dose							
adults and children 12 years and over	20 mL every 4 hours							
children 6 to under 12 years	10 mL every 4 hours							
children under 6 years	do not use							
<p>Other information</p> <ul style="list-style-type: none"> ■ each 10 mL contains: sodium 6 mg ■ store at 20-25°C (68-77°F) 								
<p>Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose</p>								
<p>Questions or comments?</p> <p>Call 1-888-594-0828 weekdays 9 AM to 5 PM EST</p>								