

Dimetapp Nighttime Cold & Cough

| | |
|--|--------------------------------------|
| Drug Facts | |
| Active ingredient (in each 10 mL) Diphenhydramine HCl, 12.5 mg.....Antihistamine/ | Purposes cough suppressant |
| Uses | |
| <ul style="list-style-type: none"> ■ temporarily relieves: <ul style="list-style-type: none"> ■ cough ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes due to hay fever (or other upper respiratory allergies) ■ relieves the impulse to cough to help you sleep | |
| Warnings | |
| Do not use | |
| <ul style="list-style-type: none"> ■ to sedate a child or to make a child sleepy ■ with any other product containing diphenhydramine, even one used on skin | |
| Ask a doctor before use if you have | |
| <ul style="list-style-type: none"> ■ trouble urinating due to an enlarged prostate gland ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema | |
| Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. | |
| When using this product | |
| <ul style="list-style-type: none"> ■ do not use more than directed ■ may cause marked drowsiness ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children | |
| Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition. | |

| Drug Facts (continued) | | | | | | | | | |
|---|---------------------|-----|------|---------------------------------------|---------------------|------------------------------|---------------------|------------------------|------------|
| <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> | | | | | | | | | |
| Directions | | | | | | | | | |
| <ul style="list-style-type: none"> ■ do not take more than 6 doses in any 24-hour period ■ measure only with dosage cup provided ■ keep dosage cup with product ■ mL = milliliter | | | | | | | | | |
| <table border="1"> <thead> <tr> <th>age</th> <th>dose</th> </tr> </thead> <tbody> <tr> <td>adults and children 12 years and over</td> <td>20 mL every 4 hours</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>10 mL every 4 hours</td> </tr> <tr> <td>children under 6 years</td> <td>do not use</td> </tr> </tbody> </table> | | age | dose | adults and children 12 years and over | 20 mL every 4 hours | children 6 to under 12 years | 10 mL every 4 hours | children under 6 years | do not use |
| age | dose | | | | | | | | |
| adults and children 12 years and over | 20 mL every 4 hours | | | | | | | | |
| children 6 to under 12 years | 10 mL every 4 hours | | | | | | | | |
| children under 6 years | do not use | | | | | | | | |
| Other information | | | | | | | | | |
| <ul style="list-style-type: none"> ■ each 10 mL contains: sodium 8 mg ■ store at 20-25°C (68-77°F) ■ do not refrigerate | | | | | | | | | |
| Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose | | | | | | | | | |
| Questions or comments? | | | | | | | | | |
| Call 1-888-594-0828 weekdays 9 AM to 5 PM EST | | | | | | | | | |

Dimetapp Cold & Cough

| | |
|---|-----------------|
| Drug Facts | |
| Active ingredients (in each 10 mL) Brompheniramine maleate, 2 mg.....Antihistamine Dextromethorphan HBr, 10 mg.....Cough suppressant | Purposes |
| Uses | |
| <ul style="list-style-type: none"> ■ temporarily relieves: <ul style="list-style-type: none"> ■ cough ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes due to hay fever (or other upper respiratory allergies) ■ temporarily alleviates the intensity of coughing | |
| Warnings | |
| Do not use | |
| <ul style="list-style-type: none"> ■ to sedate a child or to make a child sleepy ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. | |
| Ask a doctor before use if you have | |
| <ul style="list-style-type: none"> ■ trouble urinating due to an enlarged prostate gland ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema | |
| Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. | |
| When using this product | |
| <ul style="list-style-type: none"> ■ do not use more than directed ■ may cause marked drowsiness ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children | |
| Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition. | |
| If pregnant or breast-feeding, ask a health professional before use. | |
| Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. | |

| Drug Facts (continued) | | | | | | | | | |
|---|---------------------|-----|------|---------------------------------------|---------------------|------------------------------|---------------------|------------------------|------------|
| Directions | | | | | | | | | |
| <ul style="list-style-type: none"> ■ do not take more than 6 doses in any 24-hour period ■ measure only with dosage cup provided ■ keep dosage cup with product ■ mL = milliliter | | | | | | | | | |
| <table border="1"> <thead> <tr> <th>age</th> <th>dose</th> </tr> </thead> <tbody> <tr> <td>adults and children 12 years and over</td> <td>20 mL every 4 hours</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>10 mL every 4 hours</td> </tr> <tr> <td>children under 6 years</td> <td>do not use</td> </tr> </tbody> </table> | | age | dose | adults and children 12 years and over | 20 mL every 4 hours | children 6 to under 12 years | 10 mL every 4 hours | children under 6 years | do not use |
| age | dose | | | | | | | | |
| adults and children 12 years and over | 20 mL every 4 hours | | | | | | | | |
| children 6 to under 12 years | 10 mL every 4 hours | | | | | | | | |
| children under 6 years | do not use | | | | | | | | |
| Other information | | | | | | | | | |
| <ul style="list-style-type: none"> ■ each 10 mL contains: sodium 6 mg ■ store at 20-25°C (68-77°F) | | | | | | | | | |
| Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose | | | | | | | | | |
| Questions or comments? | | | | | | | | | |
| Call 1-888-594-0828 weekdays 9 AM to 5 PM EST | | | | | | | | | |