

Dimetapp Multi-Symptom Cold & Flu

Drug Facts	
Active ingredients (in each 10 mL)	Purposes
Acetaminophen, 320 mg Pain reliever/fever reducer	
Diphenhydramine HCl, 12.5 mg Antihistamine/cough suppressant	
Uses	
<ul style="list-style-type: none"> ■ temporarily relieves these common cold and flu symptoms: <ul style="list-style-type: none"> ■ minor aches and pains ■ sore throat ■ runny nose ■ sneezing ■ temporarily alleviates the intensity of coughing ■ temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if</p> <ul style="list-style-type: none"> ■ adult takes more than 6 doses in 24 hours, which is the maximum daily amount ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product. <p>Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> ■ skin reddening ■ blisters ■ rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	
Do not use	
<ul style="list-style-type: none"> ■ to sedate a child or to make a child sleepy ■ in a child under 6 years of age ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist. ■ with any other product containing diphenhydramine, even one used on skin 	

Drug Facts (continued)								
Ask a doctor before use if user has								
<ul style="list-style-type: none"> ■ liver disease ■ trouble urinating due to an enlarged prostate gland ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema 								
Ask a doctor or pharmacist before use if user is								
<ul style="list-style-type: none"> ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers 								
When using this product								
<ul style="list-style-type: none"> ■ do not use more than directed ■ may cause marked drowsiness ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children 								
Stop use and ask a doctor if								
<ul style="list-style-type: none"> ■ pain or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition. 								
If pregnant or breast-feeding, ask a health professional before use.								
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.								
Directions								
<ul style="list-style-type: none"> ■ do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage. ■ measure only with dosage cup provided ■ keep dosage cup with product ■ mL = milliliter 								
<table border="1"> <thead> <tr> <th>age</th> <th>dose</th> </tr> </thead> <tbody> <tr> <td>adults and children 12 years and over</td> <td>20 mL every 4 hours do not take more than 6 doses in any 24-hour period</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>10 mL every 4 hours do not give more than 5 doses in any 24-hour period</td> </tr> <tr> <td>children under 6 years</td> <td>do not use</td> </tr> </tbody> </table>	age	dose	adults and children 12 years and over	20 mL every 4 hours do not take more than 6 doses in any 24-hour period	children 6 to under 12 years	10 mL every 4 hours do not give more than 5 doses in any 24-hour period	children under 6 years	do not use
age	dose							
adults and children 12 years and over	20 mL every 4 hours do not take more than 6 doses in any 24-hour period							
children 6 to under 12 years	10 mL every 4 hours do not give more than 5 doses in any 24-hour period							
children under 6 years	do not use							

Drug Facts (continued)
Other information
<ul style="list-style-type: none"> ■ each 10 mL contains: sodium 6 mg ■ store at 20-25°C (68-77°F) ■ read and keep carton
Inactive ingredients
anhydrous citric acid, artificial flavor, edetate disodium, FD&C red no. 40, glycerin, menthol, polyethylene glycol, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose
Questions or comments?
Call 1-888-594-0828 weekdays 9 AM to 5 PM EST