

# Children's Dimetapp<sup>®</sup> Cold & Allergy

Brompheniramine maleate, USP 2 mg, Phenylephrine HCl, USP 5 mg

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

## **Drug Facts**

### **Active ingredients (in each 10 ml)**

### **Purposes**

Brompheniramine maleate, USP 2 mg .....	Antihistamine
Phenylephrine HCl, USP 5 mg .....	Nasal decongestant

### **Uses**

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
- temporarily restores freer breathing through the nose

### **Warnings**

#### **Do not use**

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### **Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as emphysema, asthma, or chronic bronchitis

#### **Ask a doctor or pharmacist before use if you are**

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

#### **When using this product**

- **do not use more than directed**
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

# Children's Dimetapp<sup>®</sup> Cold & Allergy

Brompheniramine maleate, USP 2 mg, Phenylephrine HCl, USP 5 mg

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

## Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosing cup with product
- ml = milliliter

age	dose
adults and children 12 years and over	20 ml every 4 hours
children 6 to under 12 years	10 ml every 4 hours
Children under 6 years	do not use

## Other information

- **each 10 ml contains:** sodium 6 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients** anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

**Questions or comments?** Call weekdays from 9 AM to 5 PM EST at 1-800-762-4675